

Projectively equivariant quantization over $\mathbb{R}^{p|q}$

Fabian Radoux

13 September 2011

Introduction

Projectively
equivariant
quantization over

$\mathbb{R}^p | \mathfrak{q}$

Fabian Radoux

- Classical observables: functions on a symplectic manifold (\mathcal{M}, ω) .

Introduction

Projectively
equivariant
quantization over

$\mathbb{R}^p | \mathfrak{q}$

Fabian Radoux

- Classical observables: functions on a symplectic manifold (\mathcal{M}, ω) .
- Quantum observables: operators on a Hilbert space \mathcal{H} .

Introduction

Projectively
equivariant
quantization over

$\mathbb{R}^p|\mathfrak{q}$

Fabian Radoux

- Classical observables: functions on a symplectic manifold (\mathcal{M}, ω) .
- Quantum observables: operators on a Hilbert space \mathcal{H} .
- Dirac problem: find a bijection $Q : \mathcal{C}^\infty(\mathcal{M}) \rightarrow \mathcal{L}(\mathcal{H})$ such that

Introduction

Projectively
equivariant
quantization over

$\mathbb{R}^p|\mathfrak{q}$

Fabian Radoux

- Classical observables: functions on a symplectic manifold (\mathcal{M}, ω) .
- Quantum observables: operators on a Hilbert space \mathcal{H} .
- Dirac problem: find a bijection $Q : \mathcal{C}^\infty(\mathcal{M}) \rightarrow \mathcal{L}(\mathcal{H})$ such that
 - $Q(\mathbf{1}_{\mathcal{M}}) = \text{Id}_{\mathcal{H}}$,

Introduction

Projectively
equivariant
quantization over

$\mathbb{R}^p|q$

Fabian Radoux

- Classical observables: functions on a symplectic manifold (\mathcal{M}, ω) .
- Quantum observables: operators on a Hilbert space \mathcal{H} .
- Dirac problem: find a bijection $Q : \mathcal{C}^\infty(\mathcal{M}) \rightarrow \mathcal{L}(\mathcal{H})$ such that
 - $Q(\mathbf{1}_{\mathcal{M}}) = \text{Id}_{\mathcal{H}}$,
 - $Q(\bar{f}) = Q(f)^*$,

Introduction

Projectively
equivariant
quantization over

$\mathbb{R}^{p|q}$

Fabian Radoux

- Classical observables: functions on a symplectic manifold (\mathcal{M}, ω) .
- Quantum observables: operators on a Hilbert space \mathcal{H} .
- Dirac problem: find a bijection $Q : \mathcal{C}^\infty(\mathcal{M}) \rightarrow \mathcal{L}(\mathcal{H})$ such that
 - $Q(\mathbf{1}_{\mathcal{M}}) = \text{Id}_{\mathcal{H}}$,
 - $Q(\bar{f}) = Q(f)^*$,
 - $Q(\{f, g\}) = \frac{i}{\hbar}[Q(f), Q(g)]$.

- Hamiltonian H , observable F ,

$$\frac{dF}{dt} = \{H, F\} \longmapsto \frac{dQ(F)}{dt} = \frac{i}{\hbar} [Q(H), Q(F)]$$

- Hamiltonian H , observable F ,

$$\frac{dF}{dt} = \{H, F\} \longmapsto \frac{dQ(F)}{dt} = \frac{i}{\hbar} [Q(H), Q(F)]$$

- **Prequantization:** if $\mathcal{H} = L^2(\mathcal{M})$, if $\omega = d\alpha$,

- Hamiltonian H , observable F ,

$$\frac{dF}{dt} = \{H, F\} \longmapsto \frac{dQ(F)}{dt} = \frac{i}{\hbar} [Q(H), Q(F)]$$

- **Prequantization:** if $\mathcal{H} = L^2(\mathcal{M})$, if $\omega = d\alpha$,

$$Q(f) = \frac{\hbar}{i} X_f + f - \langle X_f, \alpha \rangle.$$

- Problem of the prequantization: if $\mathcal{M} = T^*M$, $L^2(T^*M)$ is too big \longrightarrow reduction of \mathcal{H} .

- Problem of the prequantization: if $\mathcal{M} = T^*M$, $L^2(T^*M)$ is too big \longrightarrow reduction of \mathcal{H} .
- Polarization of (\mathcal{M}, ω) : distribution $P \subset T\mathcal{M}$.

- Problem of the prequantization: if $\mathcal{M} = T^*M$, $L^2(T^*M)$ is too big \longrightarrow reduction of \mathcal{H} .
- Polarization of (\mathcal{M}, ω) : distribution $P \subset T\mathcal{M}$.
- P allows to reduce \mathcal{H} , \mathcal{H} is replaced by \mathcal{H}^P .

- Problem of the prequantization: if $\mathcal{M} = T^*M$, $L^2(T^*M)$ is too big \longrightarrow reduction of \mathcal{H} .
- Polarization of (\mathcal{M}, ω) : distribution $P \subset T\mathcal{M}$.
- P allows to reduce \mathcal{H} , \mathcal{H} is replaced by \mathcal{H}^P .
- The observable f is quantizable if $Q(f)$ preserves \mathcal{H}^P .

- Problem of the prequantization: if $\mathcal{M} = T^*M$, $L^2(T^*M)$ is too big \longrightarrow reduction of \mathcal{H} .
- Polarization of (\mathcal{M}, ω) : distribution $P \subset T\mathcal{M}$.
- P allows to reduce \mathcal{H} , \mathcal{H} is replaced by \mathcal{H}^P .
- The observable f is quantizable if $Q(f)$ preserves \mathcal{H}^P .
- The set of quantizable observables: \mathcal{A} .

- Problem of the prequantization: if $\mathcal{M} = T^*M$, $L^2(T^*M)$ is too big \longrightarrow reduction of \mathcal{H} .
- Polarization of (\mathcal{M}, ω) : distribution $P \subset T\mathcal{M}$.
- P allows to reduce \mathcal{H} , \mathcal{H} is replaced by \mathcal{H}^P .
- The observable f is quantizable if $Q(f)$ preserves \mathcal{H}^P .
- The set of quantizable observables: \mathcal{A} .
- Geometric quantization Q_G : $Q_G = Q|_{\mathcal{A}}$.

- If $\mathcal{M} = T^*M$, if P is the vertical polarization, then

$$\mathcal{H}^P \cong L^2(M), \mathcal{A} \cong \mathcal{S}_{\leq 1}(M),$$

- If $\mathcal{M} = T^*M$, if P is the vertical polarization, then

$$\mathcal{H}^P \cong L^2(M), \mathcal{A} \cong \mathcal{S}_{\leq 1}(M),$$

$$Q_G(X^i(x)p_i + A(x)) = \frac{\hbar}{i}X^i(x)\partial_i + A(x).$$

- If $\mathcal{M} = T^*M$, if P is the vertical polarization, then

$$\mathcal{H}^P \cong L^2(M), \mathcal{A} \cong \mathcal{S}_{\leq 1}(M),$$

$$Q_G(X^i(x)p_i + A(x)) = \frac{\hbar}{i} X^i(x) \partial_i + A(x).$$

- Is it possible to extend the geometric quantization to $\mathcal{S}(M)$?

- If $\mathcal{M} = T^*M$, if P is the vertical polarization, then

$$\mathcal{H}^P \cong L^2(M), \mathcal{A} \cong \mathcal{S}_{\leq 1}(M),$$

$$Q_G(X^i(x)p_i + A(x)) = \frac{\hbar}{i}X^i(x)\partial_i + A(x).$$

- Is it possible to extend the geometric quantization to $\mathcal{S}(M)$?
- Is this prolongation unique?

- If $\mathcal{M} = T^*M$, if P is the vertical polarization, then

$$\mathcal{H}^P \cong L^2(M), \mathcal{A} \cong \mathcal{S}_{\leq 1}(M),$$

$$Q_G(X^i(x)p_i + A(x)) = \frac{\hbar}{i}X^i(x)\partial_i + A(x).$$

- Is it possible to extend the geometric quantization to $\mathcal{S}(M)$?
- Is this prolongation unique?
- Is it possible to reestablish the uniqueness?

- **Equivariant quantization:** action of a Lie group G on M :
 $\Phi : G \times M \rightarrow M$.

- **Equivariant quantization:** action of a Lie group G on M :
 $\Phi : G \times M \rightarrow M$.
- Equivariant quantization Q : linear bijection Q :
 $\mathcal{S}(M) \rightarrow \mathcal{D}(M)$ s.t. $\sigma(Q(S)) = S$ and s.t.
 $Q(\Phi_g^* S) = \Phi_g^* Q(S) \forall g \in G$.

- **Equivariant quantization:** action of a Lie group G on M :
 $\Phi : G \times M \rightarrow M$.
- Equivariant quantization Q : linear bijection Q :
 $\mathcal{S}(M) \rightarrow \mathcal{D}(M)$ s.t. $\sigma(Q(S)) = S$ and s.t.
 $Q(\Phi_g^* S) = \Phi_g^* Q(S) \forall g \in G$.
- $Q(L_h^* S) = L_h^* Q(S) \forall h \in \mathfrak{g}$.

- **Equivariant quantization:** action of a Lie group G on M :
 $\Phi : G \times M \rightarrow M$.
- Equivariant quantization Q : linear bijection Q :
 $\mathcal{S}(M) \rightarrow \mathcal{D}(M)$ s.t. $\sigma(Q(S)) = S$ and s.t.
 $Q(\Phi_g^* S) = \Phi_g^* Q(S) \forall g \in G$.
- $Q(L_{h^*} S) = L_{h^*} Q(S) \forall h \in \mathfrak{g}$.
- Idea: to take G sufficiently small to have a quantization and sufficiently big to have the uniqueness.

- **Equivariant quantization:** action of a Lie group G on M :
 $\Phi : G \times M \rightarrow M$.
- Equivariant quantization Q : linear bijection Q :
 $\mathcal{S}(M) \rightarrow \mathcal{D}(M)$ s.t. $\sigma(Q(S)) = S$ and s.t.
 $Q(\Phi_g^* S) = \Phi_g^* Q(S) \forall g \in G$.
- $Q(L_h^* S) = L_h^* Q(S) \forall h \in \mathfrak{g}$.
- Idea: to take G sufficiently small to have a quantization and sufficiently big to have the uniqueness.
- **Projective case (P. Lecomte, V. Ovsienko):**
- $PGL(m+1, \mathbb{R})$ acts on $\mathbb{R}P^m$.

- **Equivariant quantization:** action of a Lie group G on M :
 $\Phi : G \times M \rightarrow M$.
- Equivariant quantization Q : linear bijection Q :
 $\mathcal{S}(M) \rightarrow \mathcal{D}(M)$ s.t. $\sigma(Q(S)) = S$ and s.t.
 $Q(\Phi_g^* S) = \Phi_g^* Q(S) \forall g \in G$.
- $Q(L_{h^*} S) = L_{h^*} Q(S) \forall h \in \mathfrak{g}$.
- Idea: to take G sufficiently small to have a quantization and sufficiently big to have the uniqueness.
- **Projective case (P. Lecomte, V. Ovsienko):**
- $PGL(m+1, \mathbb{R})$ acts on $\mathbb{R}P^m$.
- $X \in \mathfrak{sl}(m+1, \mathbb{R}) \mapsto X^*$ vector field on $\mathbb{R}P^m$.

- **Equivariant quantization:** action of a Lie group G on M :
 $\Phi : G \times M \rightarrow M$.
- Equivariant quantization Q : linear bijection Q :
 $\mathcal{S}(M) \rightarrow \mathcal{D}(M)$ s.t. $\sigma(Q(S)) = S$ and s.t.
 $Q(\Phi_g^* S) = \Phi_g^* Q(S) \forall g \in G$.
- $Q(L_{h^*} S) = L_{h^*} Q(S) \forall h \in \mathfrak{g}$.
- Idea: to take G sufficiently small to have a quantization and sufficiently big to have the uniqueness.
- **Projective case (P. Lecomte, V. Ovsienko):**
 - $PGL(m+1, \mathbb{R})$ acts on $\mathbb{R}P^m$.
 - $X \in \mathfrak{sl}(m+1, \mathbb{R}) \mapsto X^*$ vector field on $\mathbb{R}P^m$.
 - $\exists Q : L_X Q(S) = Q(L_X S) \forall X \in \mathfrak{sl}(m+1, \mathbb{R})$.

- Conformal case (C. Duval, P. Lecomte, V. Ovsienko):
- $SO(p+1, q+1)$ acts on $S^p \times S^q$.

- Conformal case (C. Duval, P. Lecomte, V. Ovsienko):
- $SO(p+1, q+1)$ acts on $S^p \times S^q$.
- $X \in \mathfrak{so}(p+1, q+1) \mapsto X^*$ vector field on \mathbb{R}^m .

- Conformal case (C. Duval, P. Lecomte, V. Ovsienko):
- $SO(p+1, q+1)$ acts on $S^p \times S^q$.
- $X \in \mathfrak{so}(p+1, q+1) \mapsto X^*$ vector field on \mathbb{R}^m .
- $\exists Q : L_X Q(S) = Q(L_X S) \forall X \in \mathfrak{so}(p+1, q+1)$.

- Conformal case (C. Duval, P. Lecomte, V. Ovsienko):
- $SO(p+1, q+1)$ acts on $S^p \times S^q$.
- $X \in \mathfrak{so}(p+1, q+1) \mapsto X^*$ vector field on \mathbb{R}^m .
- $\exists Q : L_X Q(S) = Q(L_X S) \forall X \in \mathfrak{so}(p+1, q+1)$.
- Casimir operator method:
 - ℓ: Semi-simple Lie algebra endowed with a non degenerate Killing form K ,
 - $K(X, Y) := \text{tr}(\text{ad}(X) \circ \text{ad}(Y))$

- Conformal case (C. Duval, P. Lecomte, V. Ovsienko):
- $SO(p+1, q+1)$ acts on $S^p \times S^q$.
- $X \in \mathfrak{so}(p+1, q+1) \mapsto X^*$ vector field on \mathbb{R}^m .
- $\exists Q : L_X Q(S) = Q(L_X S) \forall X \in \mathfrak{so}(p+1, q+1)$.
- Casimir operator method:
 - ℓ: Semi-simple Lie algebra endowed with a non degenerate Killing form K ,
 $K(X, Y) := \text{tr}(\text{ad}(X) \circ \text{ad}(Y))$
- (V, β) : representation of ℓ .

- Conformal case (C. Duval, P. Lecomte, V. Ovsienko):
- $SO(p+1, q+1)$ acts on $S^p \times S^q$.
- $X \in \mathfrak{so}(p+1, q+1) \mapsto X^*$ vector field on \mathbb{R}^m .
- $\exists Q : L_X Q(S) = Q(L_X S) \forall X \in \mathfrak{so}(p+1, q+1)$.
- Casimir operator method:
 - ℓ: Semi-simple Lie algebra endowed with a non degenerate Killing form K ,
 $K(X, Y) := \text{tr}(\text{ad}(X) \circ \text{ad}(Y))$
- (V, β) : representation of ℓ .
- $(u_i : i \leq n)$: basis of ℓ ; $(u'_i : i \leq n)$: Killing-dual basis
($K(u_i, u'_j) = \delta_{i,j}$).

- Conformal case (C. Duval, P. Lecomte, V. Ovsienko):
- $SO(p+1, q+1)$ acts on $S^p \times S^q$.
- $X \in \mathfrak{so}(p+1, q+1) \mapsto X^*$ vector field on \mathbb{R}^m .
- $\exists Q : L_X Q(S) = Q(L_X S) \forall X \in \mathfrak{so}(p+1, q+1)$.
- Casimir operator method:
 - ℓ: Semi-simple Lie algebra endowed with a non degenerate Killing form K ,
 $K(X, Y) := \text{tr}(\text{ad}(X) \circ \text{ad}(Y))$
- (V, β) : representation of ℓ .
- $(u_i : i \leq n)$: basis of ℓ ; $(u'_i : i \leq n)$: Killing-dual basis
($K(u_i, u'_j) = \delta_{i,j}$).
- Casimir operator corresponding to (V, β) :

$$\sum_{i=1}^n \beta(u'_i) \beta(u_i).$$

- $(\mathcal{S}(\mathbb{R}^m), L)$ and $(\mathcal{D}(\mathbb{R}^m), \mathcal{L})$ are representations of \mathfrak{g} .

- $(\mathcal{S}(\mathbb{R}^m), L)$ and $(\mathcal{D}(\mathbb{R}^m), \mathcal{L})$ are representations of \mathfrak{g} .
- C and \mathcal{C} : Casimir operators of \mathfrak{g} on $\mathcal{S}(M)$ and $\mathcal{D}(M)$.

- $(\mathcal{S}(\mathbb{R}^m), L)$ and $(\mathcal{D}(\mathbb{R}^m), \mathcal{L})$ are representations of \mathfrak{g} .
- C and \mathcal{C} : Casimir operators of \mathfrak{g} on $\mathcal{S}(M)$ and $\mathcal{D}(M)$.
- If $C(S) = \alpha S$ and $\mathcal{L} \circ Q = Q \circ L$, then
 $\mathcal{C}(Q(S)) = \alpha Q(S)$.

- $(\mathcal{S}(\mathbb{R}^m), L)$ and $(\mathcal{D}(\mathbb{R}^m), \mathcal{L})$ are representations of \mathfrak{g} .
- C and \mathcal{C} : Casimir operators of \mathfrak{g} on $\mathcal{S}(M)$ and $\mathcal{D}(M)$.
- If $C(S) = \alpha S$ and $\mathcal{L} \circ Q = Q \circ L$, then $\mathcal{C}(Q(S)) = \alpha Q(S)$.
- In non-critical situations: if $C(S) = \alpha S$, then $\exists! Q(S)$ s.t. $\mathcal{C}(Q(S)) = \alpha Q(S)$, $\sigma(Q(S)) = S$.

- $(\mathcal{S}(\mathbb{R}^m), L)$ and $(\mathcal{D}(\mathbb{R}^m), \mathcal{L})$ are representations of \mathfrak{g} .
- C and \mathcal{C} : Casimir operators of \mathfrak{g} on $\mathcal{S}(M)$ and $\mathcal{D}(M)$.
- If $C(S) = \alpha S$ and $\mathcal{L} \circ Q = Q \circ L$, then $\mathcal{C}(Q(S)) = \alpha Q(S)$.
- In non-critical situations: if $C(S) = \alpha S$, then $\exists!$ $Q(S)$ s.t. $\mathcal{C}(Q(S)) = \alpha Q(S)$, $\sigma(Q(S)) = S$.
- In these conditions: $\mathcal{L}(Q(S)) = Q(L(S))$ because:

- $(\mathcal{S}(\mathbb{R}^m), L)$ and $(\mathcal{D}(\mathbb{R}^m), \mathcal{L})$ are representations of \mathfrak{g} .
- C and \mathcal{C} : Casimir operators of \mathfrak{g} on $\mathcal{S}(M)$ and $\mathcal{D}(M)$.
- If $C(S) = \alpha S$ and $\mathcal{L} \circ Q = Q \circ L$, then $\mathcal{C}(Q(S)) = \alpha Q(S)$.
- In non-critical situations: if $C(S) = \alpha S$, then $\exists! Q(S)$ s.t. $\mathcal{C}(Q(S)) = \alpha Q(S)$, $\sigma(Q(S)) = S$.
- In these conditions: $\mathcal{L}(Q(S)) = Q(L(S))$ because:
 - $\sigma(\mathcal{L}(Q(S))) = \sigma(Q(L(S))) = L(S)$;

- $(\mathcal{S}(\mathbb{R}^m), L)$ and $(\mathcal{D}(\mathbb{R}^m), \mathcal{L})$ are representations of \mathfrak{g} .
- C and \mathcal{C} : Casimir operators of \mathfrak{g} on $\mathcal{S}(M)$ and $\mathcal{D}(M)$.
- If $C(S) = \alpha S$ and $\mathcal{L} \circ Q = Q \circ L$, then $\mathcal{C}(Q(S)) = \alpha Q(S)$.
- In non-critical situations: if $C(S) = \alpha S$, then $\exists!$ $Q(S)$ s.t. $\mathcal{C}(Q(S)) = \alpha Q(S)$, $\sigma(Q(S)) = S$.
- In these conditions: $\mathcal{L}(Q(S)) = Q(L(S))$ because:
 - $\sigma(\mathcal{L}(Q(S))) = \sigma(Q(L(S))) = L(S)$;
 - $\mathcal{C}(Q(L(S))) = \alpha Q(L(S))$, $\mathcal{C}(\mathcal{L}(Q(S))) = \alpha \mathcal{L}(Q(S))$.

Projectively equivariant quantization on $\mathbb{R}^{p|q}$ (P. Mathonet, R.)

Projectively
equivariant
quantization over
 $\mathbb{R}^{p|q}$

Fabian Radoux

- $f \in C^{\infty p|q}$:
$$f(x^1, \dots, x^p) = \sum_{I \subseteq \{1, \dots, q\}} f_I(x^1, \dots, x^p) \theta^I,$$
$$\theta^i \theta^j = -\theta^j \theta^i$$

Projectively equivariant quantization on $\mathbb{R}^{p|q}$ (P. Mathonet, R.)

Projectively
equivariant
quantization over
 $\mathbb{R}^{p|q}$

Fabian Radoux

- $f \in C^{\infty p|q}$:
$$f(x^1, \dots, x^p) = \sum_{I \subseteq \{1, \dots, q\}} f_I(x^1, \dots, x^p) \theta^I,$$
$$\theta^i \theta^j = -\theta^j \theta^i$$
- Vector field X : superderivation of $C^{\infty p|q}$:
$$X.(fg) = (X.f)g + (-1)^{\tilde{f}\tilde{X}} f(X.g)$$

Projectively equivariant quantization on $\mathbb{R}^{p|q}$ (P. Mathonet, R.)

Projectively
equivariant
quantization over
 $\mathbb{R}^{p|q}$

Fabian Radoux

- $f \in C^{\infty p|q}$:
 $f(x^1, \dots, x^p) = \sum_{I \subseteq \{1, \dots, q\}} f_I(x^1, \dots, x^p) \theta^I,$
 $\theta^i \theta^j = -\theta^j \theta^i$
- Vector field X : superderivation of $C^{\infty p|q}$:
 $X.(fg) = (X.f)g + (-1)^{\tilde{f}\tilde{X}} f(X.g)$
- λ -density over $\mathbb{R}^{p|q}$: smooth function $f \in C^{\infty p|q}$ s.t.

$$L_X^\lambda f = X(f) + \lambda \operatorname{div}(X)f,$$

where

$$\operatorname{div}(X) = \sum_{i=1}^{p+q} (-1)^{\tilde{y}_i \tilde{X}^i} \partial_{y^i} X^i.$$

Projectively equivariant quantization on $\mathbb{R}^{p|q}$ (P. Mathonet, R.)

Projectively
equivariant
quantization over
 $\mathbb{R}^{p|q}$

Fabian Radoux

- $f \in C^{\infty p|q}$:
 $f(x^1, \dots, x^p) = \sum_{I \subseteq \{1, \dots, q\}} f_I(x^1, \dots, x^p) \theta^I,$
 $\theta^i \theta^j = -\theta^j \theta^i$
- Vector field X : superderivation of $C^{\infty p|q}$:
 $X.(fg) = (X.f)g + (-1)^{\tilde{f}\tilde{X}} f(X.g)$
- λ -density over $\mathbb{R}^{p|q}$: smooth function $f \in C^{\infty p|q}$ s.t.

$$L_X^\lambda f = X(f) + \lambda \operatorname{div}(X)f,$$

where

$$\operatorname{div}(X) = \sum_{i=1}^{p+q} (-1)^{\tilde{y}_i \tilde{X}^i} \partial_{y^i} X^i.$$

- (V, ρ) : representation of $\mathfrak{gl}(p|q)$.

Projectively equivariant quantization on $\mathbb{R}^{p|q}$ (P. Mathonet, R.)

Projectively
equivariant
quantization over
 $\mathbb{R}^{p|q}$

Fabian Radoux

- $f \in C^{\infty p|q}$:
 $f(x^1, \dots, x^p) = \sum_{I \subseteq \{1, \dots, q\}} f_I(x^1, \dots, x^p) \theta^I,$
 $\theta^i \theta^j = -\theta^j \theta^i$
- Vector field X : superderivation of $C^{\infty p|q}$:
 $X.(fg) = (X.f)g + (-1)^{\tilde{f}\tilde{X}} f(X.g)$
- λ -density over $\mathbb{R}^{p|q}$: smooth function $f \in C^{\infty p|q}$ s.t.

$$L_X^\lambda f = X(f) + \lambda \operatorname{div}(X)f,$$

where

$$\operatorname{div}(X) = \sum_{i=1}^{p+q} (-1)^{\tilde{y}_i \tilde{X}^i} \partial_{y^i} X^i.$$

- (V, ρ) : representation of $\mathfrak{gl}(p|q)$.
- Space of tensor fields of type $(V, \rho) = C^{\infty p|q} \otimes V$.

■ $L_X(f \otimes v) = X(f) \otimes v + (-1)^{\tilde{X}\tilde{f}} \sum_{ij} f J_i^j \otimes \rho(e_j^i) v,$
where $J_i^j = (-1)^{\tilde{y}^i \tilde{X} + 1} (\partial_{y^i} X^j).$

- $L_X(f \otimes v) = X(f) \otimes v + (-1)^{\tilde{X}\tilde{f}} \sum_{ij} f J_i^j \otimes \rho(e_j^i) v$,
where $J_i^j = (-1)^{\tilde{y}^i \tilde{X} + 1} (\partial_{y^i} X^j)$.
- V for densities: B^δ : 1|0-dimensional vector space spanned by one element u .

- $L_X(f \otimes v) = X(f) \otimes v + (-1)^{\tilde{X}\tilde{f}} \sum_{ij} f J_i^j \otimes \rho(e_j^i) v$,
where $J_i^j = (-1)^{\tilde{y}^i \tilde{X} + 1} (\partial_{y^i} X^j)$.
- V for densities: B^δ : 1|0-dimensional vector space spanned by one element u .
- Representation ρ of $\mathfrak{gl}(p|q)$ on B^δ : $\rho(A)u = -\delta_{\text{str}}(A)u$.

- $L_X(f \otimes v) = X(f) \otimes v + (-1)^{\tilde{X}\tilde{f}} \sum_{ij} f J_i^j \otimes \rho(e_j^i) v$,
where $J_i^j = (-1)^{\tilde{y}^i \tilde{X} + 1} (\partial_{y^i} X^j)$.
- V for densities: B^δ : 1|0-dimensional vector space spanned by one element u .
- Representation ρ of $\mathfrak{gl}(p|q)$ on B^δ : $\rho(A)u = -\delta \text{str}(A)u$.
- $S_\delta^k = C^{\infty p|q} \otimes (B^\delta \otimes S^k \mathbb{R}^{p|q})$.

- $L_X(f \otimes v) = X(f) \otimes v + (-1)^{\tilde{X}\tilde{f}} \sum_{ij} f J_i^j \otimes \rho(e_j^i) v$,
where $J_i^j = (-1)^{\tilde{y}^i \tilde{X} + 1} (\partial_{y^i} X^j)$.
- V for densities: B^δ : 1|0-dimensional vector space spanned by one element u .
- Representation ρ of $\mathfrak{gl}(p|q)$ on B^δ : $\rho(A)u = -\delta \text{str}(A)u$.
- $\mathcal{S}_\delta^k = C^{\infty p|q} \otimes (B^\delta \otimes S^k \mathbb{R}^{p|q})$.
- Differential operator $D \in \mathcal{D}_{\lambda, \mu}^k$:

$$D(f) = \sum_{|\alpha| \leq k} f_\alpha \partial_{x^1}^{\alpha_1} \cdots \partial_{x^p}^{\alpha_p} \partial_{\theta^1}^{\alpha_{p+1}} \cdots \partial_{\theta^q}^{\alpha_{p+q}} f.$$

- $L_X(f \otimes v) = X(f) \otimes v + (-1)^{\tilde{X}\tilde{f}} \sum_{ij} f J_i^j \otimes \rho(e_j^i) v$,
where $J_i^j = (-1)^{\tilde{y}^i \tilde{X} + 1} (\partial_{y^i} X^j)$.
- V for densities: B^δ : 1|0-dimensional vector space spanned by one element u .
- Representation ρ of $\mathfrak{gl}(p|q)$ on B^δ : $\rho(A)u = -\delta \text{str}(A)u$.
- $\mathcal{S}_\delta^k = C^{\infty p|q} \otimes (B^\delta \otimes S^k \mathbb{R}^{p|q})$.
- Differential operator $D \in \mathcal{D}_{\lambda, \mu}^k$:

$$D(f) = \sum_{|\alpha| \leq k} f_\alpha \partial_{x^1}^{\alpha_1} \cdots \partial_{x^p}^{\alpha_p} \partial_{\theta^1}^{\alpha_{p+1}} \cdots \partial_{\theta^q}^{\alpha_{p+q}} f.$$

- $\mathcal{L}_X D = L_X^\mu \circ D - (-1)^{\tilde{X}\tilde{D}} D \circ L_X^\lambda$.

- Space of symbols: graded space associated with $\mathcal{D}_{\lambda,\mu}$.

- Space of symbols: graded space associated with $\mathcal{D}_{\lambda,\mu}$.
- Space of symbols isomorphic to
$$\mathcal{S}_\delta = \bigoplus_{l=0}^{\infty} \mathcal{S}_\delta^l, \quad \delta = \mu - \lambda.$$

- Space of symbols: graded space associated with $\mathcal{D}_{\lambda,\mu}$.
- Space of symbols isomorphic to
$$\mathcal{S}_\delta = \bigoplus_{l=0}^{\infty} \mathcal{S}_\delta^l, \quad \delta = \mu - \lambda.$$
- Isomorphism:

$$\sigma_k : \mathcal{D}^k \rightarrow \mathcal{S}^k : D \mapsto \sum_{|\alpha|=k} f_\alpha \otimes e_1^{\alpha_1} \vee \dots \vee e_{p+q}^{\alpha_{p+q}}.$$

- Space of symbols: graded space associated with $\mathcal{D}_{\lambda,\mu}$.
- Space of symbols isomorphic to
$$\mathcal{S}_\delta = \bigoplus_{l=0}^{\infty} \mathcal{S}_\delta^l, \quad \delta = \mu - \lambda.$$
- Isomorphism:

$$\sigma_k : \mathcal{D}^k \rightarrow \mathcal{S}^k : D \mapsto \sum_{|\alpha|=k} f_\alpha \otimes e_1^{\alpha_1} \vee \dots \vee e_{p+q}^{\alpha_{p+q}}.$$

- Quantization on $\mathbb{R}^{p|q}$: linear bijection $Q : \mathcal{S}_\delta \rightarrow \mathcal{D}_{\lambda,\mu}$
s.t. $\sigma_k(Q(S)) = S$ for all $S \in \mathcal{S}_\delta^k$.

Projective superalgebra of vector fields

Projectively
equivariant
quantization over

$\mathbb{R}^{p|q}$

Fabian Radoux

- $\mathfrak{pgl}(p+1|q) = \mathfrak{gl}(p+1|q)/\mathbb{R}\text{Id} \longleftrightarrow$ subalgebra of vector fields over $\mathbb{R}^{p|q}$.

Projective superalgebra of vector fields

Projectively
equivariant
quantization over

$\mathbb{R}^{p|q}$

Fabian Radoux

- $\mathfrak{pgl}(p+1|q) = \mathfrak{gl}(p+1|q)/\mathbb{R}\text{Id} \longleftrightarrow$ subalgebra of vector fields over $\mathbb{R}^{p|q}$.
- Ω subset of \mathbb{R}^{p+1} equal to $\{(x^0, \dots, x^p) : x^0 > 0\}$.

Projective superalgebra of vector fields

Projectively
equivariant
quantization over

$\mathbb{R}^{p|q}$

Fabian Radoux

- $\mathfrak{pgl}(p+1|q) = \mathfrak{gl}(p+1|q)/\mathbb{R}\text{Id} \longleftrightarrow$ subalgebra of vector fields over $\mathbb{R}^{p|q}$.
- Ω subset of \mathbb{R}^{p+1} equal to $\{(x^0, \dots, x^p) : x^0 > 0\}$.
- $H(\Omega)$: space of restrictions of homogeneous functions over $\mathbb{R}^{p+1|q}$ to Ω .

Projective superalgebra of vector fields

Projectively
equivariant
quantization over

$\mathbb{R}^{p|q}$

Fabian Radoux

- $\mathfrak{pgl}(p+1|q) = \mathfrak{gl}(p+1|q)/\mathbb{R}\text{Id} \longleftrightarrow$ subalgebra of vector fields over $\mathbb{R}^{p|q}$.
- Ω subset of \mathbb{R}^{p+1} equal to $\{(x^0, \dots, x^p) : x^0 > 0\}$.
- $H(\Omega)$: space of restrictions of homogeneous functions over $\mathbb{R}^{p+1|q}$ to Ω .
- Bijective correspondence $i : C^\infty{}^{p|q} \rightarrow H(\Omega)$.

Projective superalgebra of vector fields

Projectively
equivariant
quantization over

$\mathbb{R}^{p|q}$

Fabian Radoux

- $\mathfrak{pgl}(p+1|q) = \mathfrak{gl}(p+1|q)/\mathbb{R}\text{Id} \longleftrightarrow$ subalgebra of vector fields over $\mathbb{R}^{p|q}$.
- Ω subset of \mathbb{R}^{p+1} equal to $\{(x^0, \dots, x^p) : x^0 > 0\}$.
- $H(\Omega)$: space of restrictions of homogeneous functions over $\mathbb{R}^{p+1|q}$ to Ω .
- Bijective correspondence $i : C^\infty \mathbb{R}^{p|q} \rightarrow H(\Omega)$.
- Homomorphism $h_{p+1,q} : \mathfrak{gl}(p+1|q) \rightarrow \text{Vect}(\mathbb{R}^{p+1|q})$.

Projective superalgebra of vector fields

Projectively
equivariant
quantization over

$\mathbb{R}^{p|q}$

Fabian Radoux

- $\mathfrak{pgl}(p+1|q) = \mathfrak{gl}(p+1|q)/\mathbb{R}\text{Id} \longleftrightarrow$ subalgebra of vector fields over $\mathbb{R}^{p|q}$.
- Ω subset of \mathbb{R}^{p+1} equal to $\{(x^0, \dots, x^p) : x^0 > 0\}$.
- $H(\Omega)$: space of restrictions of homogeneous functions over $\mathbb{R}^{p+1|q}$ to Ω .
- Bijective correspondence $i : C^\infty \mathbb{R}^{p|q} \rightarrow H(\Omega)$.
- Homomorphism $h_{p+1,q} : \mathfrak{gl}(p+1|q) \rightarrow \text{Vect}(\mathbb{R}^{p+1|q})$.
- If $A \in \mathfrak{gl}(p+1|q)$, $h_{p+1,q}(A)$ preserves $H(\Omega)$.

Projective superalgebra of vector fields

Projectively
equivariant
quantization over

$\mathbb{R}^{p|q}$

Fabian Radoux

- $\mathfrak{pgl}(p+1|q) = \mathfrak{gl}(p+1|q)/\mathbb{R}\text{Id} \longleftrightarrow$ subalgebra of vector fields over $\mathbb{R}^{p|q}$.
- Ω subset of \mathbb{R}^{p+1} equal to $\{(x^0, \dots, x^p) : x^0 > 0\}$.
- $H(\Omega)$: space of restrictions of homogeneous functions over $\mathbb{R}^{p+1|q}$ to Ω .
- Bijective correspondence $i : C^{\infty p|q} \rightarrow H(\Omega)$.
- Homomorphism $h_{p+1,q} : \mathfrak{gl}(p+1|q) \rightarrow \text{Vect}(\mathbb{R}^{p+1|q})$.
- If $A \in \mathfrak{gl}(p+1|q)$, $h_{p+1,q}(A)$ preserves $H(\Omega)$.
- $\pi(h_{p+1,q}(A)) \in \text{Vect}(\mathbb{R}^{p|q})$:
 $\pi(h_{p+1,q}(A))(f) = i^{-1} \circ h_{p+1,q}(A) \circ i(f)$, where $f \in C^{\infty p|q}$.

- $\pi \circ h_{p+1,q}(\text{Id}) = 0$, thus $\pi \circ h_{p+1,q}$ induces a homomorphism from $\mathfrak{pgl}(p+1|q)$ to $\text{Vect}(\mathbb{R}^{p|q})$.

- $\pi \circ h_{p+1,q}(\text{Id}) = 0$, thus $\pi \circ h_{p+1,q}$ induces a homomorphism from $\mathfrak{pgl}(p+1|q)$ to $\text{Vect}(\mathbb{R}^{p|q})$.
- $\mathfrak{pgl}(p+1|q) \longleftrightarrow \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1$, $\mathfrak{g}_{-1} = \mathbb{R}^{p|q}$, $\mathfrak{g}_0 = \mathfrak{gl}(p|q)$ and $\mathfrak{g}_1 = (\mathbb{R}^{p|q})^*$.

- $\pi \circ h_{p+1,q}(\text{Id}) = 0$, thus $\pi \circ h_{p+1,q}$ induces a homomorphism from $\mathfrak{pgl}(p+1|q)$ to $\text{Vect}(\mathbb{R}^{p|q})$.
- $\mathfrak{pgl}(p+1|q) \longleftrightarrow \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1$, $\mathfrak{g}_{-1} = \mathbb{R}^{p|q}$, $\mathfrak{g}_0 = \mathfrak{gl}(p|q)$ and $\mathfrak{g}_1 = (\mathbb{R}^{p|q})^*$.
- \mathfrak{g}_{-1} : constant vector fields, \mathfrak{g}_0 : linear vector fields, \mathfrak{g}_1 : quadratic vector fields.

- $\pi \circ h_{p+1,q}(\text{Id}) = 0$, thus $\pi \circ h_{p+1,q}$ induces a homomorphism from $\mathfrak{pgl}(p+1|q)$ to $\text{Vect}(\mathbb{R}^{p|q})$.
- $\mathfrak{pgl}(p+1|q) \longleftrightarrow \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1$, $\mathfrak{g}_{-1} = \mathbb{R}^{p|q}$, $\mathfrak{g}_0 = \mathfrak{gl}(p|q)$ and $\mathfrak{g}_1 = (\mathbb{R}^{p|q})^*$.
- \mathfrak{g}_{-1} : constant vector fields, \mathfrak{g}_0 : linear vector fields, \mathfrak{g}_1 : quadratic vector fields.
- **Projectively equivariant quantization**: quantization Q s.t. $\mathcal{L}_{X^h} \circ Q = Q \circ L_{X^h}$ for every $h \in \mathfrak{pgl}(p+1|q)$.

Construction of the quantization

Projectively
equivariant
quantization over

$\mathbb{R}^p | \mathfrak{q}$

Fabian Radoux

- Q_{Aff} : inverse of the total symbol map σ

Construction of the quantization

Projectively
equivariant
quantization over

$\mathbb{R}^{p|q}$

Fabian Radoux

- Q_{Aff} : inverse of the total symbol map σ
- $\mathcal{L}_X S = Q_{\text{Aff}}^{-1} \circ \mathcal{L}_X \circ Q_{\text{Aff}}(S)$, for every $S \in \mathcal{S}_\delta$ and $X \in \text{Vect}(\mathbb{R}^{p|q})$.

Construction of the quantization

Projectively
equivariant
quantization over

$\mathbb{R}^{p|q}$

Fabian Radoux

- Q_{Aff} : inverse of the total symbol map σ
- $\mathcal{L}_X S = Q_{\text{Aff}}^{-1} \circ \mathcal{L}_X \circ Q_{\text{Aff}}(S)$, for every $S \in \mathcal{S}_\delta$ and $X \in \text{Vect}(\mathbb{R}^{p|q})$.
- Projectively equivariant quantization \longleftrightarrow
 $\mathfrak{pgl}(p+1|q)$ -module isomorphism from the
representation (\mathcal{S}_δ, L) to the representation $(\mathcal{S}_\delta, \mathcal{L})$.

Construction of the quantization

Projectively
equivariant
quantization over

$\mathbb{R}^{p|q}$

Fabian Radoux

- Q_{Aff} : inverse of the total symbol map σ
- $\mathcal{L}_X S = Q_{\text{Aff}}^{-1} \circ \mathcal{L}_X \circ Q_{\text{Aff}}(S)$, for every $S \in \mathcal{S}_\delta$ and $X \in \text{Vect}(\mathbb{R}^{p|q})$.
- Projectively equivariant quantization \longleftrightarrow
 $\mathfrak{pgl}(p+1|q)$ -module isomorphism from the
representation (\mathcal{S}_δ, L) to the representation $(\mathcal{S}_\delta, \mathcal{L})$.
- Map γ :

$$\gamma : \mathfrak{g} \rightarrow \mathfrak{gl}(\mathcal{S}_\delta, \mathcal{S}_\delta) : h \mapsto \gamma(h) = \mathcal{L}_{X^h} - L_{X^h}.$$

Construction of the quantization

Projectively
equivariant
quantization over

$\mathbb{R}^{p|q}$

Fabian Radoux

- Q_{Aff} : inverse of the total symbol map σ
- $\mathcal{L}_X S = Q_{\text{Aff}}^{-1} \circ \mathcal{L}_X \circ Q_{\text{Aff}}(S)$, for every $S \in \mathcal{S}_\delta$ and $X \in \text{Vect}(\mathbb{R}^{p|q})$.
- Projectively equivariant quantization \longleftrightarrow
 $\mathfrak{pgl}(p+1|q)$ -module isomorphism from the
representation (\mathcal{S}_δ, L) to the representation $(\mathcal{S}_\delta, \mathcal{L})$.
- Map γ :

$$\gamma : \mathfrak{g} \rightarrow \mathfrak{gl}(\mathcal{S}_\delta, \mathcal{S}_\delta) : h \mapsto \gamma(h) = \mathcal{L}_{X^h} - L_{X^h}.$$

- γ vanishes on $\mathfrak{g}_{-1} \oplus \mathfrak{g}_0$, $\gamma(h) : \mathcal{S}_\delta^k \rightarrow \mathcal{S}_\delta^{k-1}$.

■ Casimir operators:

- Casimir operators:
- \mathfrak{l} : Lie superalgebra endowed with a nondegenerate even supersymmetric bilinear form K .

- Casimir operators:
- \mathfrak{l} : Lie superalgebra endowed with a nondegenerate even supersymmetric bilinear form K .
- (V, β) : representation of \mathfrak{l} .

- Casimir operators:
- \mathfrak{l} : Lie superalgebra endowed with a nondegenerate even supersymmetric bilinear form K .
- (V, β) : representation of \mathfrak{l} .
- $(u_i : i \leq n)$: homogeneous basis of \mathfrak{l} ; $(u'_i : i \leq n)$: K -dual basis ($K(u_i, u'_j) = \delta_{i,j}$).

- Casimir operators:

- \mathfrak{l} : Lie superalgebra endowed with a nondegenerate even supersymmetric bilinear form K .
- (V, β) : representation of \mathfrak{l} .
- $(u_i : i \leq n)$: homogeneous basis of \mathfrak{l} ; $(u'_i : i \leq n)$: K -dual basis ($K(u_i, u'_j) = \delta_{i,j}$).
- Casimir operator of (V, β) :

$$\sum_{i=1}^n (-1)^{\tilde{u}_i} \beta(u_i) \beta(u'_i) = \sum_{i=1}^n \beta(u'_i) \beta(u_i).$$

- If $q \neq p + 1$, $\mathfrak{pgl}(p + 1|q) \cong \mathfrak{sl}(p + 1|q)$.

- If $q \neq p + 1$, $\mathfrak{pgl}(p + 1|q) \cong \mathfrak{sl}(p + 1|q)$.
- Killing form of $\mathfrak{sl}(p + 1|q)$:

$$K(A, B) = \text{str}(ad(A)ad(B)) = 2(p + 1 - q) \text{str}(AB).$$

- If $q \neq p + 1$, $\mathfrak{pgl}(p + 1|q) \cong \mathfrak{sl}(p + 1|q)$.
- Killing form of $\mathfrak{sl}(p + 1|q)$:

$$K(A, B) = \text{str}(ad(A)ad(B)) = 2(p + 1 - q) \text{str}(AB).$$

- The Casimir operator C of $\mathfrak{pgl}(p + 1|q) \cong \mathfrak{sl}(p + 1|q)$ on (S_δ^k, L) is equal to $\alpha(k, \delta)\text{Id}$, where

$$\alpha(k, \delta) = \frac{p - q}{2} \delta^2 - \frac{2k + p - q}{2} \delta + \frac{k(k + (p - q))}{(p - q + 1)}.$$

The Casimir operator \mathcal{C} of $\mathfrak{pgl}(p+1|q) \cong \mathfrak{sl}(p+1|q)$ on $(\mathcal{S}_\delta^k, \mathcal{L})$ is equal to $C + N$, where N is defined in this way:

$$N : \mathcal{S}_\delta^k \rightarrow \mathcal{S}_\delta^{k-1} : S \mapsto 2 \sum_i \gamma(\epsilon^i) L_{X^{\epsilon_i}} S,$$

where $\epsilon^r = \frac{(-1)^{\tilde{r}}}{2(p-q+1)} \varepsilon^r$.

- A value of δ is critical if there exist $k, l \in \mathbb{N}$ with $l < k$ such that $\alpha(k, \delta) = \alpha(l, \delta)$.

- A value of δ is critical if there exist $k, l \in \mathbb{N}$ with $l < k$ such that $\alpha(k, \delta) = \alpha(l, \delta)$.
- The set of critical values:

$$\mathfrak{C} = \bigcup_{k=1}^{\infty} \mathfrak{C}_k, \quad \mathfrak{C}_k = \left\{ \frac{2k - l + p - q}{p - q + 1} : l = 1, \dots, k \right\}.$$

- A value of δ is critical if there exist $k, l \in \mathbb{N}$ with $l < k$ such that $\alpha(k, \delta) = \alpha(l, \delta)$.
- The set of critical values:

$$\mathfrak{C} = \bigcup_{k=1}^{\infty} \mathfrak{C}_k, \quad \mathfrak{C}_k = \left\{ \frac{2k - l + p - q}{p - q + 1} : l = 1, \dots, k \right\}.$$

- Remarks:

- 1 “Same” values as in the classical case: m is replaced by $p - q$.

- A value of δ is critical if there exist $k, l \in \mathbb{N}$ with $l < k$ such that $\alpha(k, \delta) = \alpha(l, \delta)$.
- The set of critical values:

$$\mathfrak{C} = \bigcup_{k=1}^{\infty} \mathfrak{C}_k, \quad \mathfrak{C}_k = \left\{ \frac{2k - l + p - q}{p - q + 1} : l = 1, \dots, k \right\}.$$

- Remarks:
 - 1 “Same” values as in the classical case: m is replaced by $p - q$.
 - 2 0 can be a critical value.

- If δ is not critical, then there exists a unique projectively equivariant quantization.

- If δ is not critical, then there exists a unique projectively equivariant quantization.
- Proof:
 - 1 For every $S \in \mathcal{S}_\delta^k$, $\exists! \hat{S}$ s.t. $\mathcal{C}(\hat{S}) = \alpha(k, \delta)\hat{S}$ and s.t. $\hat{S} = S + S_{k-1} + \cdots + S_0$, where $S_l \in \mathcal{S}_\delta^l$ for all $l \leq k-1$.

- If δ is not critical, then there exists a unique projectively equivariant quantization.
- Proof:
 - 1 For every $S \in \mathcal{S}_\delta^k$, $\exists!$ \hat{S} s.t. $\mathcal{C}(\hat{S}) = \alpha(k, \delta)\hat{S}$ and s.t.
 $\hat{S} = S + S_{k-1} + \cdots + S_0$, where $S_l \in \mathcal{S}_\delta^l$ for all $l \leq k-1$.
 - 2 $Q(S) := \hat{S}$.

- If δ is not critical, then there exists a unique projectively equivariant quantization.
- Proof:
 - 1 For every $S \in \mathcal{S}_\delta^k$, $\exists! \hat{S}$ s.t. $\mathcal{C}(\hat{S}) = \alpha(k, \delta)\hat{S}$ and s.t. $\hat{S} = S + S_{k-1} + \cdots + S_0$, where $S_l \in \mathcal{S}_\delta^l$ for all $l \leq k-1$.
 - 2 $Q(S) := \hat{S}$.
 - 3 If $S \in \mathcal{S}_\delta^k$, $Q(L_{X^h}S) = \mathcal{L}_{X^h}(Q(S))$ because they are eigenvectors of \mathcal{C} of eigenvalue $\alpha(k, \delta)$ and because their term of degree k is exactly $L_{X^h}S$.

Divergence operator:

$$\operatorname{div} : \mathcal{S}_{\delta}^k \rightarrow \mathcal{S}_{\delta}^{k-1} : S \mapsto \sum_{j=1}^{p+q} (-1)^{\tilde{y}^j} i(\varepsilon^j) \partial_{y^j} S.$$

Divergence operator:

$$\operatorname{div} : \mathcal{S}_\delta^k \rightarrow \mathcal{S}_\delta^{k-1} : S \mapsto \sum_{j=1}^{p+q} (-1)^{\tilde{y}^j} i(\varepsilon^j) \partial_{y^j} S.$$

Theorem

If δ is not critical, then the map $Q : \mathcal{S}_\delta \rightarrow \mathcal{D}_{\lambda, \mu}$ defined by

$$Q(S)(f) = \sum_{r=0}^k C_{k,r} Q_{\text{Aff}}(\operatorname{div}^r S)(f), \quad \text{for all } S \in \mathcal{S}_\delta^k$$

is the unique $\mathfrak{sl}(p+1|q)$ -equivariant quantization if

$$C_{k,r} = \frac{\prod_{j=1}^r ((p-q+1)\lambda + k - j)}{r! \prod_{j=1}^r (p-q+2k-j - (p-q+1)\delta)} \quad \text{for all } r \geq 1.$$

- **Case $q = p + 1$:** $\mathfrak{pgl}(p + 1|p + 1)$ not endowed with a non degenerate bilinear symmetric invariant form.

- **Case $q = p + 1$:** $\mathfrak{pgl}(p + 1|p + 1)$ not endowed with a non degenerate bilinear symmetric invariant form.
- $\mathfrak{pgl}(p + 1|p + 1)$ is not simple, codimension one ideal:
 $\mathfrak{psl}(p + 1|p + 1);$
 $\mathfrak{pgl}(p + 1|p + 1) = \mathfrak{psl}(p + 1|p + 1) \oplus \mathbb{R}\text{Id}.$

- **Case $q = p + 1$:** $\mathfrak{pgl}(p + 1|p + 1)$ not endowed with a non degenerate bilinear symmetric invariant form.
- $\mathfrak{pgl}(p + 1|p + 1)$ is not simple, codimension one ideal:
 $\mathfrak{psl}(p + 1|p + 1)$;
 $\mathfrak{pgl}(p + 1|p + 1) = \mathfrak{psl}(p + 1|p + 1) \oplus \mathbb{R}\text{Id}.$
- Analyse of the existence of a $\mathfrak{psl}(p + 1|p + 1)$ -equivariant quantization.

- **Case $q = p + 1$:** $\mathfrak{pgl}(p + 1|p + 1)$ not endowed with a non degenerate bilinear symmetric invariant form.
- $\mathfrak{pgl}(p + 1|p + 1)$ is not simple, codimension one ideal:
 $\mathfrak{psl}(p + 1|p + 1)$;
 $\mathfrak{pgl}(p + 1|p + 1) = \mathfrak{psl}(p + 1|p + 1) \oplus \mathbb{R}\text{Id}.$
- Analyse of the existence of a $\mathfrak{psl}(p + 1|p + 1)$ -equivariant quantization.
- Killing form of $\mathfrak{psl}(p + 1|p + 1)$ vanishes, but K defined by

$$K([A], [B]) = \text{str}AB$$

is a nondegenerate invariant supersymmetric even form.

- Casimir operator C of $\mathfrak{psl}(p+1|p+1)$ acting on \mathcal{S}_δ^k is equal to $2k(k-1)\text{Id}$.

- Casimir operator C of $\mathfrak{psl}(p+1|p+1)$ acting on \mathcal{S}_δ^k is equal to $2k(k-1)\text{Id}$.
- $\alpha_{1,\delta} = \alpha_{0,\delta}$: critical situation.

- Casimir operator C of $\mathfrak{psl}(p+1|p+1)$ acting on \mathcal{S}_δ^k is equal to $2k(k-1)\text{Id}$.
- $\alpha_{1,\delta} = \alpha_{0,\delta}$: critical situation.
- If $k \neq 1$, Q is given by the same formula as in the case $q \neq p+1$.

- Casimir operator C of $\mathfrak{psl}(p+1|p+1)$ acting on \mathcal{S}_δ^k is equal to $2k(k-1)\text{Id}$.
- $\alpha_{1,\delta} = \alpha_{0,\delta}$: critical situation.
- If $k \neq 1$, Q is given by the same formula as in the case $q \neq p+1$.
- If $k = 1$,

$$Q_1 : S \mapsto Q(S) = Q_{\text{Aff}}(S + t \operatorname{div}(S))$$

defines a $\mathfrak{psl}(p+1|p+1)$ -equivariant quantization for every $t \in \mathbb{R}$ (vector fields in $\mathfrak{psl}(p+1|p+1)$ are divergence-free).

- Casimir operator C of $\mathfrak{psl}(p+1|p+1)$ acting on \mathcal{S}_δ^k is equal to $2k(k-1)\text{Id}$.
- $\alpha_{1,\delta} = \alpha_{0,\delta}$: critical situation.
- If $k \neq 1$, Q is given by the same formula as in the case $q \neq p+1$.
- If $k = 1$,

$$Q_1 : S \mapsto Q(S) = Q_{\text{Aff}}(S + t \operatorname{div}(S))$$

defines a $\mathfrak{psl}(p+1|p+1)$ -equivariant quantization for every $t \in \mathbb{R}$ (vector fields in $\mathfrak{psl}(p+1|p+1)$ are divergence-free).

- The $\mathfrak{psl}(p+1, p+1)$ -equivariant quantizations are $\mathfrak{pgl}(p+1, p+1)$ -equivariant (equivariance with respect to the Euler vector field corresponding to Id).

- If $q = p + 1$, Q does not depend on δ and λ .